











CHICK STARTER- 10#

A COMPLETE FEED FOR STARTING AND GROWING CHICKS

As an aid in prevention of coccidiosis where immunity to coccidiosis is not desired.

ACTIVE DRUG INGREDIENT
Amprolium 0.0125% (113.7 g/ton)

GUARANTEED ANALYSIS

Crude Protein			Min.	20 %
Lysine			Min.	1.2 %
Methionine			Min.	0.5 %
Crude Fat			Min.	3.1 %
Crude Fiber			Max.	4.1 %
Calcium (Ca)	Min.	0.9	Max.	1.4 %
Phosphorous (P)			Min.	0.6 %
Salt (NaCl)	Min.	0.2	Max.	0.8 %

INGREDIENTS

Corn, Soybean Meal, Wheat Middlings, corn Distillers Dried Grains with Solubles, Calcium Carbonate, Soybean Oil, Dicalcium Phosphate, Monocalcium Phosphate, Salt, DL-Methionine, Magnesium Oxide, L-Lysine, Sodium Bicarbonate, Butylated Hydroxyanisole, Phosphoric Acid, Methylparaben, sorbic Acid, Lemon Oil, Sodium Carbonate, Ammonium Hydroxide, Calcium Hydroxide, Tertiary Butyl Hydroquinone, Propionic Acid, Propylparaben, Benzoic Acid, Amorphous Silicon Dioxide, L-Threonine, Choline Chloride, Rice Hulls, Hemicellulose Extract, Dried Bacillus subtilis Fermentation Product, Thyme White Oil, Origanum Oil, Diatomaceous Earth, Dried Bacillus amyloliquefaciens Fermentation Product, Cinnamaldehyde, Soybean Hulls, Mono- and Diglycerides of Fatty Acids, Vitamin E Supplement, Manganous Sulfate, Zinc Sulfate, Ferrous Sulfate, Yeast Culture, Hydrolyzed Yeast, Yeast Extract, Manganous Oxide, Zinc Oxide, Selenium Yeast, Niacin Supplement, Mineral Oil, Dried Trichoderma Reesei Fermentation Product, Copper Chloride, Calcium Pantothenate, Riboflavin Supplement, Vitamin A Supplement, Menadione Nicotinamide Bisulfite, Pyridoxine Hydrochloride, Selenium Selenite, Ethylenediamine Dihydroiodide, 25-Hydroxyvitamin D3, Vitamin D3 Supplement, Biotin, Thiamine Mononitrate, Vitamin B12 Supplement, Folic Acid, Ethoxyquin (a preservative).

WARNING: Use as a sole source of Amprolium.
CAUTION: Do not use Amprolium in feeds containing bentonite.
LIMITATIONS FOR USE: Feed continuously as the sole ration.

FEEDING DIRECTIONS

Feed continuously as the sole ration from hatch until laying age for pullers (approximate 16 weeks) or through the entire life for broilers. Provide plenty of fresh, clean water at all times.

WHITE, NATURES., CRUMLES
Manufactured By
Belstra Milling Co
424 15th St. SE
DeMotte, IN 46310
10 lbs. (18.18 kg.)
290004

Safety Data Sheet Available Upon Request



CHICK STARTER- 10#

A COMPLETE FEED FOR STARTING AND GROWING CHICKS

As an aid in prevention of coccidiosis where immunity to coccidiosis is not desired.

ACTIVE DRUG INGREDIENT
Amprolium 0.0125% (113.7 g/ton)

GUARANTEED ANALYSIS

Crude Protein			Min.	20 %
Lysine			Min.	1.2 %
Methionine			Min.	0.5 %
Crude Fat			Min.	3.1 %
Crude Fiber			Max.	4.1 %
Calcium (Ca)	Min.	0.9	Max.	1.4 %
Phosphorous (P)			Min.	0.6 %
Salt (NaCl)	Min.	0.2	Max.	0.8 %

INGREDIENTS

Corn, Soybean Meal, Wheat Middlings, corn Distillers Dried Grains with Solubles, Calcium Carbonate, Soybean Oil, Dicalcium Phosphate, Monocalcium Phosphate, Salt, DL-Methionine, Magnesium Oxide, L-Lysine, Sodium Bicarbonate, Butylated Hydroxyanisole, Phosphoric Acid, Methylparaben, sorbic Acid, Lemon Oil, Sodium Carbonate, Ammonium Hydroxide, Calcium Hydroxide, Tertiary Butyl Hydroquinone, Propionic Acid, Propylparaben, Benzoic Acid, Amorphous Silicon Dioxide, L-Threonine, Choline Chloride, Rice Hulls, Hemicellulose Extract, Dried Bacillus subtilis Fermentation Product, Thyme White Oil, Origanum Oil, Diatomaceous Earth, Dried Bacillus amyloliquefaciens Fermentation Product, Cinnamaldehyde, Soybean Hulls, Mono- and Diglycerides of Fatty Acids, Vitamin E Supplement, Manganous Sulfate, Zinc Sulfate, Ferrous Sulfate, Yeast Culture, Hydrolyzed Yeast, Yeast Extract, Manganous Oxide, Zinc Oxide, Selenium Yeast, Niacin Supplement, Mineral Oil, Dried Trichoderma Reesei Fermentation Product, Copper Chloride, Calcium Pantothenate, Riboflavin Supplement, Vitamin A Supplement, Menadione Nicotinamide Bisulfite, Pyridoxine Hydrochloride, Selenium Selenite, Ethylenediamine Didhydroiodide, 25-Hydroxyvitamin D3, Vitamin D3 Supplement, Biotin, Thiamine Mononitrate, Vitamin B12 Supplement, Folic Acid, Ethoxyquin (a preservative).

WARNING: Use as a sole source of Amprolium.
CAUTION: Do not use Amprolium in feeds containing bentonite.
LIMITATIONS FOR USE: Feed continuously as the sole ration.

FEEDING DIRECTIONS

Feed continuously as the sole ration from hatch until laying age for pullers (approximate 16 weeks) or through the entire life for broilers. Provide plenty of fresh, clean water at all times.

WHITE, NATURES., CRUMBLES
Manufactured By
Belstra Milling Co
424 15th St. SE
DeMotte, IN 46310
10 lbs. (18.18 kg.)

290004

Safety Data Sheet Available Upon Request

